

DR MCCRAY CLENPIQ SPLIT DOSE INSTRUCTIONS

Have you had a heart attack within the last year?	☐ Yes	□ No
Do you have any heart or blood vessel (vascular) stents?	□ Yes	□ No
Have you had any blood clots within the last year?	☐ Yes	□ No
Have you recently had a medical procedures done?	☐ Yes	□ No
Are you on any blood thinning medications such as Plavix (Clopidogrel),	☐ Yes	□ No
Coumadin (Warfarin), Lovenox (Enoxaparin), Eliquis or Pradaxa?		

7 DAYS BEFORE THE PROCEDURE

- > STOP taking iron containing products, such as a multivitamin with iron.
- > STOP eating foods with seeds or nuts, popcorn, etc.

5 DAYS BEFORE THE PROCEDURE

- > STOP taking herbal supplements, such as ginger/garlic pills, ginseng, ginkgo, etc.
- > STOP taking any non-steroidal, anti-inflammatory medication such as Ibuprofen, Motrin, Advil, Naproxen, Aleve, etc. You may continue to take Tylenol.
- Please check with the prescribing physician if on any blood thinners.
- If you are unsure about the medications you are taking, please call your prescribing Doctor's office for clarification.

1 DAY BEFORE THE PROCEDURE

- > STOP ALL SOLID FOODS. You may have only clear liquids starting when you get out of bed in the morning. Please see clear liquid auidelines.
- FOLLOW DOSING DIRECTIONS (ATTACHED)
 This solution will cause diarrhea. Your stool should continue to become lighter until it is yellow/clear in color. If you start vomiting the prep please contact our office immediately for further instructions.

THE DAY OF THE PROCEDURE

- **EAT NO FOOD.**
- Only blood pressure and heart medications may be taken the morning of the procedure with the smallest amount of water possible. Hold other medications, unless specified by your Doctor.
- FOLLOW DOSING DIRECTIONS (ATTACHED)

2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811



CLENPIQ SPLIT DOSE INSTRUCTIONS

What is CLENPIQ?

CLENPIQ is a prescription medicine that cleans your colon.

CLENPIQ is ready for you to drink right from the bottle. It does not need to be mixed or diluted.

Do not refrigerate or freeze CLENPIQ.

THE NIGHT BEFORE THE PROCEDURE AT 5:00PM

- Drink one bottle of CLENPIQ. (You may drink it straight from the bottle)
- Follow with 5 cups (8 ounces each) of clear liquid, taken at your own pace within the next 5 hours.

Between 5 - 9pm



THE DAY OF THE PROCEDURE: 5 HOURS PRIOR

- Drink the 2nd bottle of CLENPIQ. (You may drink the prep straight from the bottle.)
- Follow with 3 cups (8 ounces each) of clear liquids. Finish liquids at least 2 hours prior to colonoscopy.

5 Hours before procedure



2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811



CLEAR LIQUID GUIDELINES

START LIQUID DIET 1 DAY BEFORE YOUR PROCEDURE

BLACK COFFEE, HOT OR ICED TEA

(NO ESPRESSO, CREAM OR DAIRY)

7-UP, SPRITE, GINGER ALE

APPLE JUICE OR WHITE GRAPE JUICE

CHICKEN OR BEEF BROTH - NO CREAM OR DAIRY SOUPS

JELLO – LEMON OR LIME ONLY

POPSICLES - LEMON OR LIME ONLY

GATORADE – LEMON OR LIME OR CLEAR ONLY

WATER

ANYTHING YOU CAN SEE THROUGH SHOULD BE OKAY, AS LONG AS IT HAS NO PARTICLES. IF YOU ARE UNSURE OF SOMETHING, PLEASE FOLLOW THE ABOVE GUIDELINES OR CALL OUT OFFICE WITH QUESTIONS. MONDAY-FRIDAY 8AM-5PM (907) 276-2811

NO SOLID FOOD. NO DAIRY. NO ALCOHOL. NOTHING BLUE, PURPLE, RED OR ORANGE COLORED.

You are required to arrange a driver for your procedure. You should not work, drive or make any major decisions for up to 12 hrs after your procedure. For your safety, you cannot be released to a taxicab. To avoid cancellation, please ensure you have a reliable driver arranged prior to your procedure date.

FAILURE TO FOLLOW THESE INSTRUCTIONS COULD RESULT IN CANCELLATION OF YOUR PROCEDURE AS WELL AS A \$250.00 FEE.

Thank you for affording us the opportunity to provide you with our healthcare services.

2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811