

## DR TOMPKINS GOLYTELY ONE DAY PREP

Have you had a heart attack within the last year?	□ Yes	□ No
Do you have any heart or blood vessel (vascular) stents?	☐ Yes	□ No
Have you had any blood clots within the last year?	□ Yes	□ No
Have you recently had a medical procedure done?	☐ Yes	□ No
Are you on any blood thinning medications such as Plavix (Clopidogrel),	□ Yes	□ No
Coumadin (Warfarin), Lovenox (Enoxaparin), Eliquis or Pradaxa?		ĺ

#### 7 DAYS BEFORE THE PROCEDURE

- > STOP taking iron containing products, such as multivitamin with iron.
- > STOP taking aspirin or aspirin containing products. Tylenol is okay if needed.
- > STOP eating foods with seeds, nuts, popcorn, etc.
- > STOP eating all green leafy vegetables (spinach, kale, cabbage, lettuce, etc)
- > STOP all GLP-1 agonist medication such as Ozempic, Rybelsus, Trulicity, Mounjaro, etc. If you take this medication for diabetes, please contact your managing provider for guidance on preventing hyperglycemia.

#### **5 DAYS BEFORE THE PROCEDURE**

- > STOP taking herbal supplements, such as ginger/garlic pills, ginseng, etc.
- > STOP taking any non-steroidal, anti-inflammatory medication such as Ibuprofen, Motrin, Advil, Naproxen, Aleve, etc. You may continue to take Tylenol.
- Please check with the prescribing physician if on any blood thinners.
- If you are unsure about the medications you are taking, please call your prescribing Doctor's office for clarification.

### 1 DAY BEFORE THE PROCEDURE

- > STOP ALL SOLID FOOD. You may only have clear liquids starting when you get out of bed in the morning. Please see clear liquid guidelines.
- ➤ Mix the PEG 3350/Golytely/Nulytely according to the directions on the container and refrigerate.
- At <u>6:00pm</u> begin drinking the PEG 3350/Golytely/Nulytely at a rate of 8-10 oz glasses every 20 minutes. This will cause diarrhea. Your stool should continue to because lighter until it is yellow/clear in color. **YOU MUCH DRINK THE ENITRE JUG OF SOLUTION**.
- Continue on your clear liquid diet until midnight.

#### ON THE DAY OF YOUR PROCEDURE

- If your procedure is scheduled at 1pm or later, you can continue clear liquids up until 4 hours before your procedure time. Then, **nothing by mouth** at least 4 hours prior to your procedure.
- Only blood pressure and heart medications may be taken the morning of the procedure with the smallest amount of water possible.

2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811



# CLEAR LIQUID GUIDELINES START LIQUID DIET 1 DAY BEFORE YOUR PROCEDURE

BLACK COFFEE, HOT OR ICED TEA

(NO ESPRESSO, CREAM OR DAIRY)

7-UP, SPRITE, GINGER ALE

APPLE JUICE OR WHITE GRAPE JUICE

CHICKEN OR BEEF BROTH - NO CREAM OR DAIRY SOUPS

JELLO – LEMON OR LIME ONLY

POPSICLES – LEMON OR LIME ONLY

GATORADE - LEMON OR LIME OR CLEAR ONLY

WATER

ANYTHING YOU CAN SEE THROUGH SHOULD BE OKAY, AS LONG AS IT HAS NO PARTICLES. IF YOU ARE UNSURE OF SOMETHING, PLEASE FOLLOW THE ABOVE GUIDELINES OR CALL OUT OFFICE WITH QUESTIONS. MONDAY-FRIDAY 8AM-5PM (907) 276-2811

NO SOLID FOOD. NO DAIRY. NO ALCOHOL. NOTHING BLUE, PURPLE, RED OR ORANGE COLORED.

\*You are required to arrange a driver for your procedure. You should not work, drive or make any major decisions for up to 12 hrs after your procedure. For your safety, you cannot be released to a taxicab. To avoid cancellation, please ensure you have a reliable driver arranged prior to your procedure date.\*

FAILURE TO FOLLOW THESE INSTRUCTIONS COULD RESULT IN CANCELLATION OF YOUR PROCEDURE AS WELL AS A \$250.00 FEE.

2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811



# Thank you for affording us the opportunity to provide you with our healthcare services.

2841 Debarr Road Building A, Suite 50 Anchorage, AK 99508 Phone: (907) 276-2811